

THE UNITED SIMILES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Northrup, King & Co.

Tolhereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF SEVENTERM YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS ASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

SOYBEAN

'S 1244'

In Testimonn Minercot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington this 28th day of January in the year of our Lord one thousand nine hundred and seventy-seven

Commissioner
Plant Variety Protection Office
Grain Division

Secretary of Agriculture

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION HYATTSVILLE, MARYLAND 20782

APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE

INSTRUCTIONS: See Reverse,				
1. VARIETY NAME OR TEMPORARY DESIGNATION	2. KIND NAME			AL USE ONLY
S 1244	Soybeans		75000 86	
3. GENUS AND SPECIES NAME	4. FAMILY NAME (Botanical)		FILING DATE	TIME
	Leguminosae		5. 8. 75 FEE RECEIVED	2:30 P.M.
Glycine max (L.) Merr.	S. DATE OF DETER		\$ 250.00	\$
	\$250,00		\$	
	January 1	970	\$250,00	\$
6. NAME OF APPLICANT(S)	7. ADDRESS (Street a	8. TELEPHONE AREA CODE AND NUMBER		
Northrup, King & Co.	P. 0. Box	959		
		is, Minnesota	55440	612-781-8011
9. IF THE NAMED APPLICANT IS NOT A PE ORGANIZATION: (Corporation, partnership,		10. STATE OF INCOM	RPORATION	11. DATE OF INCOR- PORATION
Corporation	•	Minnesota		1896
12. Name and mailing address of applic	ant representative(s	s), if any, to serve	in this application as	-
Northrup, King & Co. P. O. Box 959 Minneapolis, Minnesota	55440		·	
13. CHECK BOX BELOW FOR EACH ATTACH		,		
X 13A. Exhibit A, Origin and Bree	eding History of the	Variety (See Section	n 52 of the Plant Va	riety Protection Act.)
🛣 138. Exhibit B, Botanical Desc	ription of the Varie	ty		•
13C. Exhibit C, Objective Desc.	ription of the Variet	y		
🛣 13D. Exhibit D, Data Indicative	of Novelty			
X 13E. Exhibit E, Statement of the	e Basis of Applican	t's Ownership		
14A. Does the applicant(s) specify that (See Section 83(a), (If "Yes," an	seed of this variety	y be sold by variety elow.)	name only as a clas	ss of certified seed?
148. Does the applicant(s) specify that				rations of production
limited as to number of generation	ns?	beyond breed foundatio		CERTIFIED
The applicant declares that a viable s ance of a certificate and will be reple				
The undersigned applicant(s) of this uniform, and stable as required in Se Plant Variety Protection Act.				
Applicant is informed that false repr	esentation herein ca	an jeopardize protec	tion and result in pe	nalties.
March 1975		(800°	al Ru	0.1
(DATE)		(SI	GNATURE OF APPLICA	NT)
V			() 00	001
(DATE)	_	(si	GNATURE OF APPLICA	NT)

INSTRUCTIONS

GENERAL: Send an original copy of the application, exhibits and \$250.00 fee to U.S. Dept. of Agriculture, Agricultural Marketing Service, Grain Division, 6525 Belcrest Road, Hyattsville, Maryland 20782. (See Section 180.175 of the regulations and rules of practice.) Retain one copy for your files. All items on the face of the form are self-explanatory unless noted below.

ITEM

- Insert the date the applicant determined that he had a new variety based on the definition in Section 41 (a) of the Act and decision is made to increase the seed.
- 13a First, give the genealogy, including public and commercial MAY varieties, lines, or clones used, and the breeding method Second, give the details of subsequent stages of selection and multiplication. Third, indicate the type and frequency of variants during reproduction and multiplication and state how these variants may be identified. Fourth, provide evidence on stability.
- 13b First, give any special characteristics of the seed and of his the plant as it passes through the seedling stage, flowering stage and the fruiting stage. Second, describe the mature plant and compare it with a similar commercial variety grown under the same conditions, and indicate the differences.
- 13c A supplemental form will be furnished by the PVPO to describe in detail a variety for each kind of seed.
- 13d Provide complete data indicative of novelty. Seed and plant specimens or photographs of seed and plant comparisons clearly indicating novelty may be submitted. Seeds submitted may be sterile.
- 13e Indicate whether applicant is the actual breeder, the employer of the breeder, the owner through purchase or inheritance, etc.



EXHIBIT A ORIGIN AND BREEDING HISTORY OF S 1244 SOYBEANS

- 1969 60 plants were selected from an F_{λ} bulk population from the cross 'Corsoy' x 'Wayne'. The population had been advanced to the F_{λ} generation by harvesting 2 pods from approximately 400 plants in each generation.
- 1969-70 Seeds from each selected plant were grown in a progeny row. One of these was designated 9414. Each row was bulk-harvested if uniform.
- 1970 9414 was yield tested at Hudson, Iowa. On the basis of its high yield and maturity, it was chosen as an experimental variety worthy of further testing.
- 1971 9414 was yield tested at Stanton, Minnesota; Homer, Michigan; and Hudson, Iowa. Also, 100 representative plants were harvested individually to be grown as progeny rows in 1972.
- 1972 9414 was yield tested at Stanton, Minnesota; Delavan, Wisconsin; and Dixon, Illinois. One hundred progeny rows were grown, and any rows containing off-type plants were discarded. The rest were bulk-harvested to produce pedigree seed of the variety.
- 1973 9414 was yield tested at Stanton and Minnesota Lake, Minnesota; and Hudson, Dayton, and Washington, Iowa. An increase block was planted from the pedigree seed produced in 1972 and was harvested to produce breeder seed.
- 9414 was tested at all the locations listed for 1973 plus Darien,
 Wisconsin and Van Wert, Ohio. It was tested in University trials in
 Iowa, Minnesota, and Wisconsin. A further seed increase was made
 from the breeder seed produced in 1973. In addition, pedigree seed
 was produced as in 1972. The pedigree method of maintaining varietal
 purity will continue as long as the variety is produced.
- 9414 was named S 1244 and released to foundation seed producers.
 S 1244 is stable and uniform for all normal descriptive characteristics. A very low frequency of variants would be expected through
 mutation, outcrossing, or mechanical mixture. These will be prevented from becoming a significant constituent of the variety through
 application of the time-proven pedigree method referred to above.



EXHIBIT B BOTANICAL DESCRIPTION OF S 1244 SOYBEANS

I. Seed and Seedling.

Cotyledons of S 1244 are yellow. Seeds have dull yellow seed coats and black hyla. Seeds are fairly large (16.2 grams per 100 vs. 16.0 for Steele, 14.0 for Chippewa 64, and 15.2 for Hodgson averaged over 6 tests). Seed shape is spherical, or similar to most common varieties. Seedling hypocotyl color is purple.

II. Flowering.

When planted about May 15, S 1244 will begin flowering in about 42 days at Washington, Iowa; about the same as for Chippewa 64. Duration of flowering is similar to Chippewa 64, and flowering pattern is similar to other indeterminate, Maturity Group I varieties. Flower color is purple.

III. Fruiting.

Flowering and beginning pod set overlap, as is true of other indeterminate varieties. At full vegetative growth, S 1244 has medium-sized, ovate leaflets which are a medium green color. Canopy type is intermediate between Hark, which is narrow and open, and Chippewa 64, which is bushy and closed.

IV. Disease Reaction.

S 1244 is similar to most northern soybean varieties in its susceptibility to common foliar diseases. It is susceptible to Phytophthora root rot.

V. Mature Plant.

S 1244 has tawny or brown pubescence and brown pods. Plant height is very similar to Chippewa 64 and Steele. Lodging resistance is similar to these varieties. Most pods are 2 or 3 seeded, and there are normally several pods per node, depending upon yield level. In 15 NK trials, S 1244 has an average yield of about 110% of Chippewa 64. It is about 2 days later than Chippewa 64.

EXHIBIT D

DATA INDICATIVE OF NOVELTY FOR S 1244 SOYBEANS

S 1244 is most similar in appearance and plant habit to Chippewa 64 of the commonly grown maturity Group I varieties. It is susceptible to Phytophthora root rot; Chippewa 64 is resistant. S 1244 has brown pubescence in contrast to the grey pubescent varieties Hark, Steele, Hodgson, Peterson 85, Blackhawk, SRF 150, and A 100. S 1244 can be differentiated from Osage by hilum color (black vs. yellow). S 1244 has normal brown pubescence vs. light brown pubescence for Dunn. S 1244 cannot be differentiated from the less common maturity Group I varieties Chippewa, Wirth, Rampage, and Anoka by pubescence, flower, or hilum color or Phytophthora reaction. S 1244 is significantly later than Chippewa 64, Chippewa, Anoka, or Wirth. It has dull seed coat luster, vs. shiny luster for Chippewa, Chippewa 64, Rampage, Wirth, and Anoka.

Data Contrasting S 1244 from other maturity Group I varieties.

	Co	olor of:			Seed Coat
	Pubescence	Flower	Hi lum	Phytophthora	luster
s 1244	В	P :	в1	S	D
Chippewa	В	P	В1	S	S
Chippewa 64	В	P	в1	R	S
Hark	G	P	Y	S	D
Steele	G	P	Y	R	D
Hodgson	G	P	BF	S	S
Wirth	В	P	в1	S	S
Osage	В	P	Y	s	-
Rampage	В	P	ві	S	S
Anoka	В	P	в1	s	S
Dunn	LB	P	в1	s	s
Peterson 85	G	P	BF	s	-
Blackhawk	G	W	IB	R	s
SRF 150	G	P	Y	s	-
A 100	G	W	BF	s	s

EXHIBIT D DATA INDICATIVE OF NOVELTY FOR S 1244 SOYBEANS

I. Seed.

A. Seed Description.

Seed of S 1244 have dull yellow seed coats, yellow cotyledons, and black hyla, and are spherical in shape.

B. Seed Size.

Variety	Wt. in g per 100 seeds*
	<u> </u>
S 1244 Steele Chippewa 64 Hodgson Corsoy	16.2 16.0 14.0 15.2 13.8

^{*} Average of 6 trials.

C. Chemical Composition of Seed.

<u>Variety</u>	Protein %	<u>0il %</u>
S 1244	41.4	19.6
Chippewa 64	38.6	20.1
Hark	40.0	20.0
Corsoy	37.6	20.8

II. Leaf and Canopy Characteristics.

S 1244 has medium-sized ovate leaflets which are a medium green color. Leaf size and canopy type are intermediate between Hark, which has small leaflets and a narrow, open canopy, and Chippewa 64, which has larger leaflets and a more bushy, closed canopy.

III. Flower Golor is Burple.



IV. Mature Plant.

- A. Description. S 1244 has tawny pubescence and brown pods.
- B. Agronomic Data.

Variety	<u>Y</u> ield (K/ha) (X ₁₅ 1972-74)	Maturity (X ₃ 1973-74)	_Lodging* (X ₁₁ 1973-74)	Height (cm) (X ₂ 1973-74)
S 1244 Chippewa 64	2603 2366	√9 – 14 9 – 12	1.7	76 78
Anoka Steele Hark	23 86 2 7 52	9 - 13 9-15 9-21	2.0 1.9 1.8	66 76 83

^{* 1 =} erect; 5 = prostrate.

V. Susceptible to Phytophthora root rot.

INSTRUCTIONS: See Reverse.

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION HYATTSVILLE, MARYLAND 20782 OBJECTIVE DESCRIPTION OF VARIETY SOYBEAN (GLYCINE MAX)

EXHIBIT C (Soybean)

NAME OF APPLICANT(S)		FOR OFF	ICIAL USE ONLY
Northrup, King & Co. ADDRESS (Street and No., or R.F.D. No.; City, State, and ZIP Code)		PVPO NUMBER	7500086
P O Pau Of O	Ì	VARIETY NAME OR	· · · · · · · · · · · · · · · · · · ·
P. O. Box 959 Minneapolis, Minnesota 55440		S 1244	·
		• •	<u> </u>
Place the appropriate number that describes the varietal char-	acter of this variety	y in the boxes be	low.
1 1 = SPHERICAL 2 = SPHERICAL 3 = ELONGAT	E 4 = OTHER (Specify)	
2. SEED COAT COLOR:	ı,	SHADE:	
1 = YELLOW 2 = GREEN 3 = BROWN 5 = OTHER (Specify)	4 = BLACK	2 1 = LIGHT	2 = MEDIUM 3 = DARK
3. SEED COAT LUSTER:	4. SEED SIZE		·
1 = DULL 2 = SHINY	1 6 GRAMS PER	100 SEEDS	
5. HILUM COLOR:	1	SHADE:	
6 1 = BUFF 2 = YELLOW 3 = BROWN 4 = GRAY	5 ± IMPERFECT		
6 = BLACK 7 = OTHER (Specify)	BLACK	3 1 = LIGHT	2 = MEDIUM 3 = DARK
6. COTYLEDON COLOR:	7. LEAFLET SIZE (Se	ee Reverse):	
1 = YELLOY 2 = GREEN	2 1 = SMALL	2 = MEDIUM	3 = LARGE
8. LEAFLET SHAPE:			
1 = OVATE 2 = OBLONG 3 = LANCEOLATE 4 =	ELLIPTICAL 5 = 0	OTHER (Specify)	
9. LEAF COLOR (See reverse):		10. FLOWER COLO	₹:
2 = MEDIUM GREEN 3 = DARK	GREEN	1 = WHITE 3 = OTHER (2 = PURPLE
11. POD COLOR:	12: POD SET:	<u> </u>	
2 1 = TAN 2 = BROWN 3 = BLACK	2 1 = SCATTER	ED 2 = CONCE	ENTRATED
13. PLANT PUBESCENCE COLOR:	1	SHADE:	
2 = BROWN 3 = OTHER (Specity)	1	2 1 = LIGHT	2 = MEDIUM 3 = DARK
14. PLANT TYPES (See Reverse):	15. PLANT HABIT:	<u> </u>	· ·,• ·, · · -·
3 1 = SLENDER 2 = BUSHY 3 = INTERMEDIATE	2 1 = DETERMIN 3 = OTHER (S)		TERMINATE
16. HYPOCOTYL COLOR:	17. SEED PROTEIN:		
2 T = GREEN 2 = PURPLE	1 = A 2	:	ੂ 'ਚੋਂ -
18. NUMBER OF DAYS TO FLOWERING 19. MATURITY GROUP: (Place a zero in first box (e.g. 0 9) when 1 = 00 2	=0 3 = 1	4 = 11	5 = m·
days are 9 or less.) 4 2 3 6 = IV 7	= v 8 = vı	9 = VII	10 = VIII
20. SIZE OF 10 DAY OLD SEEDLING GROWN UNDER CONSTANT LIGHT	(Growth Chamber) AT		
(e.g. [0 2]) when size is 9 mm. or loss.) MM. LENGTH MM. LENGTH		MM. WIDTH	
21. DISEASE: (Enter 0 =Not Tested; 1 = Susceptible; 2 = Resistant)		OF COTYLEDO	······································
BACTERIAL O SOYBEAN O DOWNY	PURPLE 1	POD AND STEM BLIGHT	O ROOT
O FROGEYE 1 STEM 1 PHYTO- 1	BROWN 0	TARGET	BROWN
BUD RHIZOCTONIA	OTHER (Specify)	sрот (L'_ spot 10004
			



EXHIBIT E STATEMENT OF THE BASIS OF APPLICANT'S OWNERSHIP

The soybean variety, S 1244, was developed by Northrup, King & Co.'s breeding staff at its Washington, Iowa research farm from germ plasm sources cited in Exhibit A of this application. Northrup, King & Co. believes that the variety it has created is novel as defined in the Plant Variety Protection Act and, therefore, that Northrup, King & Co. is the sole owner of the variety.

CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant shape	Corsoy	Petiole angle	Corsoy
Leaf shape	Corsoy	Seed size	Steele
Leaf color	Chippewa 64	Seed shape	Steel e
Leaf surface	Corsoy	Seedling pigmentation	Chippewa 64

23. GIVE DATA FOR SU				1	SIZE	CON	TENT	AVERAGE NO.	······
VARIETY	NO. OF DAYS	SCORE	PLANT HEIGHT	Width	Length	Protein	Oil	OF PODS PER PLANT	IODINE NO.
Submitted	122	1.7	76 cm	50 mm	94 mm	41.4	19.6 %	18 @ 350000plt/h	1
Name of similar variety Chippews 64.	120	1.6	78 cm	57 mm	88 mm	38.6	20.1	19 @ 350000p1t/b	

INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for completing this form:

- 1. Scott, Walter O. and Samuel R. Aldrich, 1970, Modern Soybean Production, The Farmer Quarterly.
- 2. Norman, A. G., 1963, The Soybean: Genetics, Breeding, Physiology, Nutrition, Management.
- 3. McKie, J. W., and K. L. Anderson, 1970, The Soybean Book.

L'EAF COLOR: Nickerson's or any recognized color fan may be used to determine the leaf color of the described variety. The following Soybean varieties may be used as a guide to identify the colors listed on the form.

COLOR	VARIETY
Light Green	''Ada''
Medium Green	''Wilkin''
Dark Green	''Swift''

LEAF SIZE: The following varieties may be used as a guide to identify the relative size leaves.

SIZE	VARIETY
Small	''Amsoy''
Medium	"Bonus"
Large	''Anoka''

PLANT TYPE: The following varieties may be used as a guide to identify the plant type



